



#9

Patent  
MBHB00,801-F 400.003

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re the Application of:

Blatt, et al.

Serial No.: 09/740,332

Filed: December 18, 2000

For: ENZYMATIC NUCLEIC ACID  
TREATMENT OF DISEASES OR  
CONDITIONS RELATED TO HEPATITIS  
C VIRUS INFECTION

)  
) **Group Art Unit:**  
)  
) **Examiner:** not assigned  
)  
)  
)  
)  
)  
)  
)

PRELIMINARY AMENDMENT

Commissioner for Patents  
Washington, D.C. 20231

Sir:

Applicant respectfully request that this amendment be entered prior to examination of the above-mentioned application on the merits. It is believed that no fee is due for filing this response; however, if a fee is due, the Commissioner is authorized to charge our Deposit Account No. 132490.

IN THE SPECIFICATION:


On page 1, after the first paragraph, please insert the following new paragraph:

" The Sequence Listing file named "MBHB00-801-F Sequence Listing.txt" submitted on Compact Disc-Recordable (CD-R) medium ("010326\_2122") in compliance with 37 C.F.R. §1.52(e) is incorporated herein by reference."

CERTIFICATE OF MAILING (37 C.F.R. 1.8a)

I hereby certify that this correspondence is being deposited with the United States Postal Service as first class mail in an envelope addressed to the: Assistant Commissioner for Patents, Washington D.C. 20231, on March 26, 2001.

Date: March 7, 2001

  
Anita J. Terpstra

On page 15, please replace paragraph 2 with the following substitute paragraph:

" By "consists essentially of" is meant that the active enzymatic nucleic acid molecule of the invention contains an enzymatic center or core equivalent to those in the examples, and binding arms able to bind RNA such that cleavage at the target site occurs. Other sequences may be present which do not interfere with such cleavage. Thus, a core region may, for example, include one or more loop or stem-loop structures, which do not prevent enzymatic activity. Such sequences can be designated as "X", for example, as in a loop or stem/loop structure. For example, a core sequence for a hammerhead enzymatic nucleic acid can be 5'- CUGAUGAG -3' and 5' - CGAA -3' connected by "X", where X= 5' - GCCGUUAGGC -3' (SEQ ID NO: 9704), or any other stem II region known in the art. Similarly, for other enzymatic nucleic acid molecules of the instant invention, additional sequences may be present that do not interfere with the function of the nucleic acid molecule."

Please replace originally-filed pages 69-72 with the substitute pages 69-72 submitted herewith.

**IN THE ABSTRACT**

On page 176, please replace the originally-filed abstract with the following substitute abstract:

"The present invention relates to enzymatic nucleic acid molecules (e.g., ribozymes and DNazymes) that modulate the expression and/or replication of hepatitis C virus (HCV)."

**IN THE FIGURES:**

Please replace Figures 1-5, 7 and 8 with substitute Figures 1-5, 7 and 8, submitted herewith.

**SEQUENCE LISTING:**

Please incorporate by reference the sequence listing submitted herewith. The sequences in the sequence listing are found in the originally-filed specification and figures.

**REMARKS**

**Specification**

Page 1 of the specification was amended to incorporate by reference the sequence listing, submitted herewith.

Page 15, paragraph 2, of the specification was amended to provide clarification and to include the appropriate SEQ ID NO, in compliance with 37 C.F.R § 1.821-1.825.

Pages 69-72 of the specification were amended to provide the appropriate reference numbers for the references cited in Table 1.

No new matter has been added by way of any of these amendments to the specification.

A copy of the marked up original sheets for the described pages are attached as Appendix A and Appendix B.

**Abstract**

The abstract was amended merely to provide clarification. A copy of the marked up original abstract is attached as Appendix C.

**Figures**

New Figures 1-5, 7 and 8 were amended to include the appropriate SEQ ID NOs in compliance with 37 C.F.R § 1.821-1.825. No new matter has been added by way of these amendments.

A marked-up version of the original Figures 1-5, 7 and 8 are attached as Appendix D.


**Sequence Listing**

A sequence listing is attached which includes the sequences found in the specification and drawings. In compliance with 37 C.F.R § 1.821-1.825 and § 1.52(e), the applicant herewith submits the Sequence Listing on Compact Disc-Recordable (CD-R) medium in duplicate (COPY 1 and COPY 2), in lieu of the paper copy under 37 C.F.R. § 1.821(c), and substitute computer readable form copy (COPY 3). The paper and computer readable forms of the Sequence Listing are the same. The Statements under 37 CFR § 1.821(f) and § 1.52(e) are also provided.

The Sequence Listing has been generated from the specification and does not constitute new subject matter. The Sequence Listing has been prepared in the PatentIn Ver.2.0 format and checked with Checker Version 3.0 Program. No error has been found.

Respectfully submitted,

**McDonnell Boehnen Hulbert & Berghoff**

  
Anita J. Terpstra  
Registration No. 47,132

McDONNELL BOEHNEN  
HULBERT & BERGHOFF  
300 South Wacker Drive  
Chicago, Illinois 60606  
312/913-0001 (telephone)  
312/913-0002 (facsimile)

## Appendix A

By “consists essentially of” is meant that the active enzymatic nucleic acid molecule of the invention contains an enzymatic center or core equivalent to those in the examples, and binding arms able to bind RNA such that cleavage at the target site occurs. Other sequences may be present which do not interfere with such cleavage. Thus, a core region may, for example, include one or more loop or stem-loop structures, which do not prevent enzymatic activity. Such sequences can be designated as “X”, for example, as in a loop or stem/loop structure. For example, a [A] core sequence for a hammerhead enzymatic nucleic acid can be 5'- CUGAUGAG -3' [X] and 5' - CGAA -3' connected by "X", where X= 5' - GCCGUUAGGC -3' (SEQ ID NO: 9704), or any other stem II region known in the art. Similarly, for other enzymatic nucleic acid molecules of the instant invention, additional sequences may be present that do not interfere with the function of the nucleic acid molecule.

## Appendix B

### Hepatitis Delta Virus (HDV) Ribozyme

- Size: ~60 nucleotides.
- Trans cleavage of target RNAs demonstrated [<sup>xxxix</sup>].
- Binding sites and structural requirements not fully determined, although no sequences 5' of cleavage site are required. Folded ribozyme contains a pseudoknot structure [<sup>xl</sup>].
- Reaction mechanism: attack by 2'-OH 5' to the scissile bond to generate cleavage products with 2',3'-cyclic phosphate and 5'-OH ends.
- Only 2 known members of this class. Found in human HDV.
- Circular form of HDV is active and shows increased nuclease stability [<sup>xli</sup>]

i Michel, Francois; Westhof, Eric. Slippery substrates. *Nat. Struct. Biol.* (1994), 1(1), 5-7.

ii Lisacek, Frederique; Diaz, Yolande; Michel, Francois. Automatic identification of group I intron cores in genomic DNA sequences. *J. Mol. Biol.* (1994), 235(4), 1206-17.

iii Herschlag, Daniel; Cech, Thomas R.. Catalysis of RNA cleavage by the *Tetrahymena thermophila* ribozyme. 1. Kinetic description of the reaction of an RNA substrate complementary to the active site. *Biochemistry* (1990), 29(44), 10159-71.

iv Herschlag, Daniel; Cech, Thomas R.. Catalysis of RNA cleavage by the *Tetrahymena thermophila* ribozyme. 2. Kinetic description of the reaction of an RNA substrate that forms a mismatch at the active site. *Biochemistry* (1990), 29(44), 10172-80.

v Knitt, Deborah S.; Herschlag, Daniel. pH Dependencies of the *Tetrahymena* Ribozyme Reveal an Unconventional Origin of an Apparent pKa. *Biochemistry* (1996), 35(5), 1560-70.

vi Bevilacqua, Philip C.; Sugimoto, Naoki; Turner, Douglas H.. A mechanistic framework for the second step of splicing catalyzed by the *Tetrahymena* ribozyme. *Biochemistry* (1996), 35(2), 648-58.

vii Li, Yi; Bevilacqua, Philip C.; Mathews, David; Turner, Douglas H.. Thermodynamic and activation parameters for binding of a pyrene-labeled substrate by the *Tetrahymena* ribozyme: docking is not diffusion-controlled and is driven by a favorable entropy change. *Biochemistry* (1995), 34(44), 14394-9.

viii Banerjee, Alope Raj; Turner, Douglas H.. The time dependence of chemical modification reveals slow steps in the folding of a group I ribozyme. *Biochemistry* (1995), 34(19), 6504-12.

ix Zarrinkar, Patrick P.; Williamson, James R.. The P9.1-P9.2 peripheral extension helps guide folding of the *Tetrahymena* ribozyme. *Nucleic Acids Res.* (1996), 24(5), 854-8.

- x Strobel, Scott A.; Cech, Thomas R.. Minor groove recognition of the conserved G.cntdot.U pair at the Tetrahymena ribozyme reaction site. *Science* (Washington, D. C.) (1995), 267(5198), 675-9.
- xi Strobel, Scott A.; Cech, Thomas R.. Exocyclic Amine of the Conserved G.cntdot.U Pair at the Cleavage Site of the Tetrahymena Ribozyme Contributes to 5'-Splice Site Selection and Transition State Stabilization. *Biochemistry* (1996), 35(4), 1201-11.
- xii Sullenger, Bruce A.; Cech, Thomas R.. Ribozyme-mediated repair of defective mRNA by targeted trans-splicing. *Nature* (London) (1994), 371(6498), 619-22.
- xiii Robertson, H.D.; Altman, S.; Smith, J.D. *J. Biol. Chem.*, 247, 5243-5251 (1972).
- xiv Forster, Anthony C.; Altman, Sidney. External guide sequences for an RNA enzyme. *Science* (Washington, D. C., 1883-) (1990), 249(4970), 783-6.
- xv Yuan, Y.; Hwang, E. S.; Altman, S. Targeted cleavage of mRNA by human RNase P. *Proc. Natl. Acad. Sci. USA* (1992) 89, 8006-10.
- xvi Harris, Michael E.; Pace, Norman R.. Identification of phosphates involved in catalysis by the ribozyme RNase P RNA. *RNA* (1995), 1(2), 210-18.
- xvii Pan, Tao; Loria, Andrew; Zhong, Kun. Probing of tertiary interactions in RNA: 2'-hydroxyl-base contacts between the RNase P RNA and pre-tRNA. *Proc. Natl. Acad. Sci. U. S. A.* (1995), 92(26), 12510-14.
- xviii Pyle, Anna Marie; Green, Justin B.. Building a Kinetic Framework for Group II Intron Ribozyme Activity: Quantitation of Interdomain Binding and Reaction Rate. *Biochemistry* (1994), 33(9), 2716-25.
- xix Michels, William J. Jr.; Pyle, Anna Marie. Conversion of a Group II Intron into a New Multiple-Turnover Ribozyme that Selectively Cleaves Oligonucleotides: Elucidation of Reaction Mechanism and Structure/Function Relationships. *Biochemistry* (1995), 34(9), 2965-77.
- xx Zimmerly, Steven; Guo, Huatao; Eskes, Robert; Yang, Jian; Perlman, Philip S.; Lambowitz, Alan M.. A group II intron RNA is a catalytic component of a DNA endonuclease involved in intron mobility. *Cell* (Cambridge, Mass.) (1995), 83(4), 529-38.
- xxi Griffin, Edmund A., Jr.; Qin, Zhifeng; Michels, Williams J., Jr.; Pyle, Anna Marie. Group II intron ribozymes that cleave DNA and RNA linkages with similar efficiency, and lack contacts with substrate 2'-hydroxyl groups. *Chem. Biol.* (1995), 2(11), 761-70.
- xxii Michel, Francois; Ferat, Jean Luc. Structure and activities of group II introns. *Annu. Rev. Biochem.* (1995), 64, 435-61.
- xxiii Abramovitz, Dana L.; Friedman, Richard A.; Pyle, Anna Marie. Catalytic role of 2'-

hydroxyl groups within a group II intron active site. Science (Washington, D. C.) (1996), 271(5254), 1410-13.

xxiv Daniels, Danette L.; Michels, William J., Jr.; Pyle, Anna Marie. Two competing pathways for self-splicing by group II introns: a quantitative analysis of in vitro reaction rates and products. J. Mol. Biol. (1996), 256(1), 31-49.

xxv Guo, Hans C. T.; Collins, Richard A.. Efficient trans-cleavage of a stem-loop RNA substrate by a ribozyme derived from Neurospora VS RNA. EMBO J. (1995), 14(2), 368-76.

xxvi Scott, W.G., Finch, J.T., Aaron, K. The crystal structure of an all RNA hammerhead ribozyme: A proposed mechanism for RNA catalytic cleavage. Cell, (1995), 81, 991-1002.

xxvii McKay, Structure and function of the hammerhead ribozyme: an unfinished story. RNA, (1996), 2, 395-403.

xxviii Long, D., Uhlenbeck, O., Hertel, K. Ligation with hammerhead ribozymes. US Patent No. 5,633,133.

xxix Hertel, K.J., Herschlag, D., Uhlenbeck, O. A kinetic and thermodynamic framework for the hammerhead ribozyme reaction. Biochemistry, (1994) 33, 3374-3385. Beigelman, L., *et al.*, Chemical modifications of hammerhead ribozymes. J. Biol. Chem., (1995) 270, 25702-25708.

xxx Beigelman, L., *et al.*, Chemical modifications of hammerhead ribozymes. J. Biol. Chem., (1995) 270, 25702-25708.

xxxi Hampel, Arnold; Tritz, Richard; Hicks, Margaret; Cruz, Phillip. 'Hairpin' catalytic RNA model: evidence for helices and sequence requirement for substrate RNA. Nucleic Acids Res. (1990), 18(2), 299-304.

xxxii Chowrira, Bharat M.; Berzal-Herranz, Alfredo; Burke, John M.. Novel guanosine requirement for catalysis by the hairpin ribozyme. Nature (London) (1991), 354(6351), 320-2.

xxxiii Berzal-Herranz, Alfredo; Joseph, Simpson; Chowrira, Bharat M.; Butcher, Samuel E.; Burke, John M.. Essential nucleotide sequences and secondary structure elements of the hairpin ribozyme. EMBO J. (1993), 12(6), 2567-73.

xxxiv Joseph, Simpson; Berzal-Herranz, Alfredo; Chowrira, Bharat M.; Butcher, Samuel E.. Substrate selection rules for the hairpin ribozyme determined by in vitro selection, mutation, and analysis of mismatched substrates. Genes Dev. (1993), 7(1), 130-8.

xxxv Berzal-Herranz, Alfredo; Joseph, Simpson; Burke, John M.. In vitro selection of active hairpin ribozymes by sequential RNA-catalyzed cleavage and ligation reactions. Genes Dev. (1992), 6(1), 129-34.

xxxvi Hegg, Lisa A.; Fedor, Martha J.. Kinetics and Thermodynamics of Intermolecular Catalysis by Hairpin Ribozymes. Biochemistry (1995), 34(48), 15813-28.



xxxvii Grasby, Jane A.; Mersmann, Karin; Singh, Mohinder; Gait, Michael J.. Purine Functional Groups in Essential Residues of the Hairpin Ribozyme Required for Catalytic Cleavage of RNA. *Biochemistry* (1995), 34(12), 4068-76.

xxxviii Schmidt, Sabine; Beigelman, Leonid; Karpeisky, Alexander; Usman, Nassim; Sorensen, Ulrik S.; Gait, Michael J.. Base and sugar requirements for RNA cleavage of essential nucleoside residues in internal loop B of the hairpin ribozyme: implications for secondary structure. *Nucleic Acids Res.* (1996), 24(4), 573-81.

xxxix Perrotta, Anne T.; Been, Michael D.. Cleavage of oligoribonucleotides by a ribozyme derived from the hepatitis .delta. virus RNA sequence. *Biochemistry* (1992), 31(1), 16-21.

xl Perrotta, Anne T.; Been, Michael D.. A pseudoknot-like structure required for efficient self-cleavage of hepatitis delta virus RNA. *Nature (London)* (1991), 350(6317), 434-6.


xli Puttaraju, M.; Perrotta, Anne T.; Been, Michael D.. A circular trans-acting hepatitis delta virus ribozyme. *Nucleic Acids Res.* (1993), 21(18), 4253


## Appendix C

### Abstract of the Disclosure

The present invention relates to [E]enzymatic nucleic acid molecules (e.g., ribozymes and DNAzymes) that modulate the expression and/or replication of hepatitis C virus (HCV).

**Figure 1: Examples of Chemically Stabilized Enzymatic Nucleic Acid motifs**






All ribo G's in the Zinzyme can be replaced with 2'-O-methyl G, or combinations thereof




Figure 5: DNAzyme Motif



Legend


Y = U or C  
R = A or G

**Figure 7: Secondary structure of the HCV 5'UTR**





**Figure 8: A Chemically Stabilized Enzymatic Nucleic Acid Molecule**



UPPER CASE = RIBO nucleotide

lower case = 2'-O-methyl nucleotide

u = 2'-deoxy-2'-amino Uridine

s = phosphorothioate

B = inverted deoxybasic moiety

## Hepatitis Delta Virus (HDV) Ribozyme

- Size: ~60 nucleotides.
- Trans cleavage of target RNAs demonstrated [<sup>xxxix</sup>].
- 5 • Binding sites and structural requirements not fully determined, although no sequences 5' of cleavage site are required. Folded ribozyme contains a pseudoknot structure [<sup>xi</sup>].
- Reaction mechanism: attack by 2'-OH 5' to the scissile bond to generate cleavage products with 2',3'-cyclic phosphate and 5'-OH ends.
- Only 2 known members of this class. Found in human HDV.
- 10 • Circular form of HDV is active and shows increased nuclease stability [<sup>xli</sup>]
- i Michel, Francois; Westhof, Eric. Slippery substrates. *Nat. Struct. Biol.* (1994), 1(1), 5-7.
- ii Lisacek, Frederique; Diaz, Yolande; Michel, Francois. Automatic identification of group I intron cores in genomic DNA sequences. *J. Mol. Biol.* (1994), 235(4), 1206-17.
- 15 iii Herschlag, Daniel; Cech, Thomas R.. Catalysis of RNA cleavage by the *Tetrahymena thermophila* ribozyme. 1. Kinetic description of the reaction of an RNA substrate complementary to the active site. *Biochemistry* (1990), 29(44), 10159-71.
- 20 iv Herschlag, Daniel; Cech, Thomas R.. Catalysis of RNA cleavage by the *Tetrahymena thermophila* ribozyme. 2. Kinetic description of the reaction of an RNA substrate that forms a mismatch at the active site. *Biochemistry* (1990), 29(44), 10172-80.
- 25 v Knitt, Deborah S.; Herschlag, Daniel. pH Dependencies of the *Tetrahymena* Ribozyme Reveal an Unconventional Origin of an Apparent pKa. *Biochemistry* (1996), 35(5), 1560-70.
- 30 vi Bevilacqua, Philip C.; Sugimoto, Naoki; Turner, Douglas H.. A mechanistic framework for the second step of splicing catalyzed by the *Tetrahymena* ribozyme. *Biochemistry* (1996), 35(2), 648-58.
- 35 vii Li, Yi; Bevilacqua, Philip C.; Mathews, David; Turner, Douglas H.. Thermodynamic and activation parameters for binding of a pyrene-labeled substrate by the *Tetrahymena* ribozyme: docking is not diffusion-controlled and is driven by a favorable entropy change. *Biochemistry* (1995), 34(44), 14394-9.
- 40 viii Banerjee, Alope Raj; Turner, Douglas H.. The time dependence of chemical modification reveals slow steps in the folding of a group I ribozyme. *Biochemistry* (1995), 34(19), 6504-12.
- ix Zarrinkar, Patrick P.; Williamson, James R.. The P9.1-P9.2 peripheral extension helps guide folding of the *Tetrahymena* ribozyme. *Nucleic Acids Res.* (1996), 24(5), 854-8.

- x Strobel, Scott A.; Cech, Thomas R.. Minor groove recognition of the conserved G.cntdot.U pair at the Tetrahymena ribozyme reaction site. *Science* (Washington, D. C.) (1995), 267(5198), 675-9.
- xi Strobel, Scott A.; Cech, Thomas R.. Exocyclic Amine of the Conserved G.cntdot.U Pair at the Cleavage Site of the Tetrahymena Ribozyme Contributes to 5'-Splice Site Selection and Transition State Stabilization. *Biochemistry* (1996), 35(4), 1201-11.
- 10 xii Sullenger, Bruce A.; Cech, Thomas R.. Ribozyme-mediated repair of defective mRNA by targeted trans-splicing. *Nature* (London) (1994), 371(6498), 619-22.
- xiii Robertson, H.D.; Altman, S.; Smith, J.D. *J. Biol. Chem.*, 247, 5243-5251 (1972).
- 15 xiv Forster, Anthony C.; Altman, Sidney. External guide sequences for an RNA enzyme. *Science* (Washington, D. C., 1883-) (1990), 249(4970), 783-6.
- xv Yuan, Y.; Hwang, E. S.; Altman, S. Targeted cleavage of mRNA by human RNase P. *Proc. Natl. Acad. Sci. USA* (1992) 89, 8006-10.
- 20 xvi Harris, Michael E.; Pace, Norman R.. Identification of phosphates involved in catalysis by the ribozyme RNase P RNA. *RNA* (1995), 1(2), 210-18.
- 25 xvii Pan, Tao; Loria, Andrew; Zhong, Kun. Probing of tertiary interactions in RNA: 2'-hydroxyl-base contacts between the RNase P RNA and pre-tRNA. *Proc. Natl. Acad. Sci. U. S. A.* (1995), 92(26), 12510-14.
- 30 xviii Pyle, Anna Marie; Green, Justin B.. Building a Kinetic Framework for Group II Intron Ribozyme Activity: Quantitation of Interdomain Binding and Reaction Rate. *Biochemistry* (1994), 33(9), 2716-25.
- 35 xix Michels, William J. Jr.; Pyle, Anna Marie. Conversion of a Group II Intron into a New Multiple-Turnover Ribozyme that Selectively Cleaves Oligonucleotides: Elucidation of Reaction Mechanism and Structure/Function Relationships. *Biochemistry* (1995), 34(9), 2965-77.
- 40 xx Zimmerly, Steven; Guo, Huatao; Eskes, Robert; Yang, Jian; Perlman, Philip S.; Lambowitz, Alan M.. A group II intron RNA is a catalytic component of a DNA endonuclease involved in intron mobility. *Cell* (Cambridge, Mass.) (1995), 83(4), 529-38.
- xxi Griffin, Edmund A., Jr.; Qin, Zhifeng; Michels, Williams J., Jr.; Pyle, Anna Marie. Group II intron ribozymes that cleave DNA and RNA linkages with similar efficiency, and lack contacts with substrate 2'-hydroxyl groups. *Chem. Biol.* (1995), 2(11), 761-70.
- 45 xxii Michel, Francois; Ferat, Jean Luc. Structure and activities of group II introns. *Annu. Rev. Biochem.* (1995), 64, 435-61.
- xxiii Abramovitz, Dana L.; Friedman, Richard A.; Pyle, Anna Marie. Catalytic role of 2'-

hydroxyl groups within a group II intron active site. *Science* (Washington, D. C.) (1996), 271(5254), 1410-13.

xxiv Daniels, Danette L.; Michels, William J., Jr.; Pyle, Anna Marie. Two competing pathways for self-splicing by group II introns: a quantitative analysis of in vitro reaction rates and products. *J. Mol. Biol.* (1996), 256(1), 31-49.

xxv Guo, Hans C. T.; Collins, Richard A.. Efficient trans-cleavage of a stem-loop RNA substrate by a ribozyme derived from *Neurospora* VS RNA. *EMBO J.* (1995), 14(2), 368-76.

xxvi Scott, W.G., Finch, J.T., Aaron, K. The crystal structure of an all RNA hammerhead ribozyme: A proposed mechanism for RNA catalytic cleavage. *Cell*, (1995), 81, 991-1002.

xxvii McKay, Structure and function of the hammerhead ribozyme: an unfinished story. *RNA*, (1996), 2, 395-403.

xxviii Long, D., Uhlenbeck, O., Hertel, K. Ligation with hammerhead ribozymes. US Patent No. 5,633,133.

xxix Hertel, K.J., Herschlag, D., Uhlenbeck, O. A kinetic and thermodynamic framework for the hammerhead ribozyme reaction. *Biochemistry*, (1994) 33, 3374-3385. Beigelman, L., *et al.*, Chemical modifications of hammerhead ribozymes. *J. Biol. Chem.*, (1995) 270, 25702-25708.

xxx Beigelman, L., *et al.*, Chemical modifications of hammerhead ribozymes. *J. Biol. Chem.*, (1995) 270, 25702-25708.

xxxi Hampel, Arnold; Tritz, Richard; Hicks, Margaret; Cruz, Phillip. 'Hairpin' catalytic RNA model: evidence for helices and sequence requirement for substrate RNA. *Nucleic Acids Res.* (1990), 18(2), 299-304.

xxxii Chowrira, Bharat M.; Berzal-Herranz, Alfredo; Burke, John M.. Novel guanosine requirement for catalysis by the hairpin ribozyme. *Nature* (London) (1991), 354(6351), 320-2.

xxxiii Berzal-Herranz, Alfredo; Joseph, Simpson; Chowrira, Bharat M.; Butcher, Samuel E.; Burke, John M.. Essential nucleotide sequences and secondary structure elements of the hairpin ribozyme. *EMBO J.* (1993), 12(6), 2567-73.

xxxiv Joseph, Simpson; Berzal-Herranz, Alfredo; Chowrira, Bharat M.; Butcher, Samuel E.. Substrate selection rules for the hairpin ribozyme determined by in vitro selection, mutation, and analysis of mismatched substrates. *Genes Dev.* (1993), 7(1), 130-8.

xxxv Berzal-Herranz, Alfredo; Joseph, Simpson; Burke, John M.. In vitro selection of active hairpin ribozymes by sequential RNA-catalyzed cleavage and ligation reactions. *Genes Dev.* (1992), 6(1), 129-34.

xxxvi Hegg, Lisa A.; Fedor, Martha J.. Kinetics and Thermodynamics of Intermolecular Catalysis by Hairpin Ribozymes. *Biochemistry* (1995), 34(48), 15813-28.

xxxvii Grasby, Jane A.; Mersmann, Karin; Singh, Mohinder; Gait, Michael J.. Purine Functional Groups in Essential Residues of the Hairpin Ribozyme Required for Catalytic Cleavage of RNA. *Biochemistry* (1995), 34(12), 4068-76.

5

xxxviii Schmidt, Sabine; Beigelman, Leonid; Karpeisky, Alexander; Usman, Nassim; Sorensen, Ulrik S.; Gait, Michael J.. Base and sugar requirements for RNA cleavage of essential nucleoside residues in internal loop B of the hairpin ribozyme: implications for secondary structure. *Nucleic Acids Res.* (1996), 24(4), 573-81.

10


xxxix Perrotta, Anne T.; Been, Michael D.. Cleavage of oligoribonucleotides by a ribozyme derived from the hepatitis .delta. virus RNA sequence. *Biochemistry* (1992), 31(1), 16-21.

15


xl Perrotta, Anne T.; Been, Michael D.. A pseudoknot-like structure required for efficient self-cleavage of hepatitis delta virus RNA. *Nature (London)* (1991), 350(6317), 434-6.

xli Puttaraju, M.; Perrotta, Anne T.; Been, Michael D.. A circular trans-acting hepatitis delta virus ribozyme. *Nucleic Acids Res.* (1993), 21(18), 4253


**Figure 1: Examples of Chemically Stabilized Enzymatic Nucleic Acid motifs**



**Figure 2: G-cleaver Motif**



**Figure 3: Chemically Stabilized Zinzyme Motif**



**Legend**

Uppercase indicates natural ribo residues

**C** indicates 2'-deoxy-2'-amino cytidine

Lowercase: 2'-O- Me

Subscript <sub>s</sub> indicates phosphothioate linkage

iB: 3'-3' abasic moiety

Y = U or C


G' can be G, ca, or caa

The gaaa tetraloop can be replaced by 18 atom polyethylene glycol (Spacer) or equivalent

All ribo G's in the Zinzyme can be replaced with 2'-O-methyl G, or combinations thereof




**Figure 4. Chemically Stabilized Amberzyme Motif**



The gaa loops may be replaced with loops of differing nucleic acid compositions, or with  $\epsilon$  linker, for example an 18 atom polyethylene glycol (Spacer) or equivalent.

Phosphorothioate linkages can be introduced, for example, at the 4 terminal 5'-internucleotide linkages, to increase nuclease stability.

Figure 5: DNAzyme Motif



Legend

Y = U or C  
R = A or G

**Figure 7: Secondary structure of the HCV 5'UTR**

